

Paleo-European crust of the Italian Western Alps: Geological history of the Argentera Massif and comparison with Mont Blanc-Aiguilles Rouges and Maures-Tanneron Massifs

Roberto Compagnoni, Simona Ferrando, Bruno Lombardo, Nicholas Radulesco, Daniela Rubatto

Journal of the Virtual Explorer, Electronic Edition, ISSN 1441-8142, volume 36, paper 3In: (Eds.) Marco Beltrando, Angelo Peccerillo, Massimo Mattei, Sandro Conticelli, and Carlo Doglioni, The Geology of Italy: tectonics and life along plate margins, 2010.

Download from: http://virtualexplorer.com.au/article/2010/228/argentera-mont-blanc-maures

Click http://virtualexplorer.com.au/subscribe/ to subscribe to the Journal of the Virtual Explorer. Email team@virtualexplorer.com.au to contact a member of the Virtual Explorer team.

Copyright is shared by The Virtual Explorer Pty Ltd with authors of individual contributions. Individual authors may use a single figure and/or a table and/or a brief paragraph or two of text in a subsequent work, provided this work is of a scientific nature, and intended for use in a learned journal, book or other peer reviewed publication. Copies of this article may be made in unlimited numbers for use in a classroom, to further education and science. The Virtual Explorer Pty Ltd is a scientific publisher and intends that appropriate professional standards be met in any of its publications.



Paleo-European crust of the Italian Western Alps: Geological history of the Argentera Massif and comparison with Mont Blanc-Aiguilles Rouges and Maures-Tanneron Massifs

Roberto Compagnoni

University of Torino - Via Valperga Caluso 35 - 10125 Torino, Italy Email: roberto.compagnoni@unito.it

Simona Ferrando

University of Torino - Via Valperga Caluso 35 - 10125 Torino, Italy Email: simona.ferrando@unito.it

Bruno Lombardo

C.N.R. Institute of Geosciences and Georesources – Via Valperga Caluso 35 – 10125 Torino, Italy *Email: bruno.lombardo@unito.it*

Nicholas Radulesco

University of Monaco - Avenue Prince Albert II - 98000 Principality of Monaco Email: nradulesco@monaco.edu

Daniela Rubatto

The Australian National University - Mills Road - 0200, Canberra Act, Australia Email: daniela.rubatto@anu.edu.au

Abstract: The "External Crystalline Massifs" of the Western Alps (Mont Blanc, Aiguilles Rouges, Grandes Rousses, Belledonne, Pelvoux, and Argentera) consist of a polymetamorphic Variscan basement, which was only marginally reworked during the Alpine tectonometamorphic cycle. These massifs experienced an early subduction event at peak metamorphic conditions of ~700°C and 1.5 GPa, followed by continental collision coupled with amphibolite-facies metamorphism, anatexis and exhumation to shallow crustal levels in the Carboniferous (see von Raumer *et al.*, 1999 for a review).

This contribution focuses on the magmatic and metamorphic history of the Argentera Massif, the southernmost and largest of the External Crystalline Massifs exposed in Italy. Its evolution from Ordovician to Early Permian is compared to that recorded in the Mont Blanc-Aiguilles Rouges Massif, the other External Crystalline Massif extensively exposed in the Italian Alps. Further comparison is drawn between these large massifs and the Maures-Tanneron Massif of Provence, France, the area of Variscan Europe nearest to Argentera.





Introduction

The Variscan orogeny (~ 380–300 Ma) is the geological event most largely represented in the basement of the European continent. It was assembled between Ordovician and Carboniferous from the larger collision of Gondwana with the northern plate of Laurentia–Baltica, which involved the microplates of Avalonia and Armorica (Matte, 2001). Variscan units (Fig. 1) extend from the Ibero-Armorican block of southern Spain to the Bohemian Massif of Poland. Large remnants of Variscan basement are preserved in the southern Variscides, within the Alpine chain, where they are located in external positions (Fig. 1).

Figure 1. European Variscides

In the Western Alps, such remnants are identified as External Crystalline Massifs (Mont Blanc, Aiguilles Rouges, Grandes Rousses, Belledonne, Pelvoux, and Argentera), which record the general evolution common to all Pangean Europe (von Raumer *et al.*, 2009). They are generally composed of a complex metamorphic basement intruded by Permo-Carboniferous granitoids. In these Massifs, the Alpine metamorphic overprint is weak and commonly limited to shear zones. The exhumation of the External Crystalline Massifs from below their sedimentary cover units and the Alpine nappes initiated in the Miocene (Argentera: Bigot-Cormier *et al.*, 2006), i.e. at the end of the Alpine orogeny.



Distribution of the European Variscan massifs and units (modified from O'Brien, 2000; Stampfli et al., 2002; von Raumer and Bussy, 2004). A: Argentera; BA: Mont Blanc – Aiguilles Rouges; C: Corsica; MT: Maures-Tanneron; RH: Rheno-Hercynian; S: Sardinia; SW: Schwarzwald; V: Vosges.

Argentera Massif

The Argentera Massif, the southernmost of the External Crystalline Massifs, straddles the boundary between Italy and France in the Maritime Alps of NW Italy and is largely composed of Variscan migmatites with abundant relicts of pre-anatectic rock types. It is an elliptical area of 60x30 km, trending WNW-ESE, with rugged relief and mountain tops of 2800-3000 m in the NW and 3100-3300 m in the centre (Figs. 2a and b). The Argentera Massif consists of the Gesso-Stura-Vésubie (GSV) Terrane to the NE and of the Tinée Terrane to the SW, which are separated by the Ferriere-Mollières shear zone (Fig. 3). The two terranes are characterized by distinct lithological associations and metamorphic evolutions, but both contain rare relicts of high-pressure (HP) and/or high-temperature (HT) mineral assemblages, which are commonly preserved within mafic rocks and exceptionally within metapelites.

The Virtual

Explorer

Modern geological maps of the Argentera Massif (Faure-Muret, 1955; Malaroda *et al.*, 1970; Malaroda, 1999) provide accurate lithological information, but fail to describe a coherent lithostratigraphy for the GSV Terrane. Such maps still use the now abandoned terminology of metamorphic rocks and migmatites of the French metasomatic school (Jung & Roques, 1952).

Gesso-Stura-Vésubie Terrane

This part of the Massif corresponds to the Malinvern-Argentera Complex of Malaroda *et al.* (1970) and comprises the Malinvern-Argentera and Chastillon-Valmasque complexes of Faure-Muret (1955).

The GSV Terrane (Fig. 3) mainly consists of migmatitic granitic gneiss—the "anatexites" of Malaroda & Schiavinato (1958) and Malaroda *et al.* (1970)—and of migmatitic paragneisses. Field relationships at the outcrop scale (Figs. 4a and b) suggest that at least a part of the "anatexites" were intrusive into the paragneiss sequence. The metasediments now appear as km-long bands in the granite gneiss, for example in the Rio Freddo valley (Bortolami & Sacchi,1968), but the intrusive contacts have been transposed by the Variscan foliation. Small bodies and boudins of mafic (amphibolite, eclogite, granulite) and ultramafic rocks are common in the "anatexites", and are more rarely exposed in the paragneisses of the GSV Terrane. Intercalations of marble and calc-silicate rocks are rare, and may contain wollastonite in addition to diopside and grossular (Calisesi, 1971). Dacitic to rhyodacitic metavolcanic rocks with transitional contacts to the surrounding "granitoid" migmatites have been locally recognized in the central and southern GSV Terrane (e.g., Rubatto et al., 2001). The eastern GSV Terrane is characterized by the Bousset-Valmasque Complex (Rubatto et al., 2001), which consists of a close association of amphibolites, agmatites with amphibolite fragments, and migmatites with amphibole-bearing anatectic granite. Carboniferous to Permian granitoids are present in the GSV Terrane, the largest pluton being the Central Granite (Fig. 3).

In the following, we briefly describe, from the oldest to the youngest, the most significant lithologies of the GSV Terrane, and summarize the information available on its geological history.

Figure 2. Argentera Massif: panoramic views





Examples of young morphology on old rocks of the GSV Terrane. (a) The W face of the Argentera ridge, looking SE from the path to Colle di Valmiana, upper Gesso della Valletta valley. (b) The E side of the Argentera ridge and the Chiotas Reservoir from the path to Rifugio Genova, looking W. Upper Valle della Rovina.



http://virtualexplorer.com.au/

Figure 3. Argentera Massif: geotectonic map



Geotectonic map of the Argentera Massif. The map is based on those of Faure-Muret (1955), Malaroda et.al. (1970), Malaroda (1999) and on sheet Gap of the 1:250'000 Geological Map of France. CG: Central Granite; VSL: Valle Stura leucogranite; FCSM: Fremamorta-Colle del Sabbione mylonite.







Migmatitic gneisses. (a) Lithologic contact between migmatitic granitic gneiss (lighter rocks and scree on the left) and migmatitic paragneisses (reddish rocks and scree on the right). GSV Terrane. Looking N from Lago del Claus, upper Gesso della Valletta valley. (b) Detail of the contact between migmatitic granitic gneiss (on the left) and migmatitic paragneiss (on the right). GSV Terrane. Laghi della Sella, upper Valle della Meris.

Relicts of HP mafic rocks

The most significant mafic bodies of the GSV Terrane are exposed in three localities (Fig. 3): 1) Laghi del Frisson, where a varied mafic sequence mainly consists of HP granulites (Figs. 5a and b); 2) Val Meris, where a large body of eclogite and retrogressed eclogite is surrounded by "anatexites" (Colombo, 1996; Rubatto et al., 2001); 3) Lago di Nasta, where amphibolitized eclogite and possibly HP granulite are both exposed (Fig. 5c). U-Pb dating on zircons suggests protolith ages for the mafic intrusions ranging between 459 ± 4 Ma (Meris eclogite, Rubatto et al., 2001) and 486 ± 7 Ma (Laghi del Frisson, Rubatto et al., 2010). Ordovician (480-460 Ma) mafic rocks, usually overprinted by low-P metamorphism, are disseminated within the External Crystalline Massifs (Ménot & Paquette, 1993; Rubatto et al., 2001; Guillot & Ménot, 2009) and, as in the Argentera, occur as relatively small bodies within the migmatitic basement.

The Laghi del Frisson mafic sequence (Fig. 5a), exposed in a tectonic window within the Helvetic-Dauphinois sedimentary cover (Fig. 3), forms a lens-shaped body, about 200 m thick and 500 m long, surrounded by biotite-bearing migmatites of broadly granitic composition ("biotite anatexite" of Malaroda *et al.*, 1970). First mapped and described as "garnet–pyroxene gneiss" by Campanino (1962 in Malaroda *et al.*, 1970), the sequence is characterized by a compositional layering that mainly consists of alternating mm- to dm-thick mafic layers of

green-brownish Pl-poor and gray-whitish Pl-rich HP granulite. The rocks of the sequence are medium- to finegrained, do not show evidence of partial melting, and are characterized by a pervasive mylonitic foliation parallel to the compositional layering (Colombo *et al.*, 1994; Ferrando *et al.*, 2008; Fig. 5b). The sequence shows a high degree of crustal contamination (Rubatto *et al.*, 2010).

This chemical signature is common to other Ordovician mafic rocks exposed in the other External Crystalline Massifs, though these are often associated with Sirich magmas, which are lacking in the Laghi del Frisson sequence. The systematic and repeated interlayering of the two main compositional types and their geochemical characteristics are inherited from a magmatic protolith, most likely a mafic layered intrusion. The original igneous layers were strongly reworked and folded by a pervasive mylonitic deformation, locally observed all through the GSV Terrane and predating widespread migmatization in the Late Carboniferous (see below). Lack of migmatisation of the mafic sequence is attributed to its more refractory composition when compared to the surrounding migmatites (Ferrando *et al.*, 2008).



Figure 5. GSV Terrane: HP mafic rocks



(a) Lago Frisson and the ridge between the upper Val Grande di Vernante and Val Gesso. The banded HP granulites are exposed in the small hill by the lake and in the ridge to the right of the saddle (Passo della Mena).
 (b) Lago Frisson HP granulites. Typical compositional banding parallel to a pervasive mylonitic foliation. Banding consists of grey-whitish PI-rich HP granulite and reddish PI-poor HP granulite.
 (c) Banded amphibolite with portions containing remnants of eclogitic garnet. GSV Terrane. Lago di Nasta, upper Gesso della Valletta valley.

Both the Pl-poor and Pl-rich HP granulites contain several generations of minerals which, coupled with thermobarometric data, define four metamorphic stages (Fig. 6; Ferrando et al., 2008). The HP granulite-facies peak (stage A: 735 \pm 15 °C, ~ 1.38 GPa) is recorded in the core of porphyroclastic garnet and omphacite in association with plagioclase, rutile \pm amphibole \pm quartz. The first decompression (stage B: ~ 710 °C and 1.10 GPa) corresponds to the rim of porphyroclastic garnet and omphacite in equilibrium with a second generation of plagioclase, rutile ± amphibole ± quartz. Mylonitization (stage C) is marked by neoblastic garnet, diopside, plagioclase, titanite \pm amphibole \pm quartz, and occurred at amphibolite-facies conditions, i.e. pressures of 0.85 GPa and still relatively HT (665±15 °C). Finally, during stage D (500 < T < 625 °C; P < 0.59 GPa) plagioclase + amphibole symplectites replaced the rim of garnet and clinopyroxene. The evolution and peak metamorphic conditions recorded by the Laghi del Frisson mafic sequence are similar to those recorded by the Meris eclogite ($T \sim$ 750°C and $P \sim 1.5$ GPa; Colombo, 1996; Rubatto et al., 2001). Based on the common metamorphic conditions, Ferrando et al. (2008) concluded that the Laghi del Frisson HP granulites and the Meris eclogite underwent the same metamorphic cycle and that the two rock types preserve different peak assemblages only because of their different bulk chemical composition.

Geochronology of HP rocks from the Argentera was first attempted by Paquette et al. (1989) using the zircon isotope-dilution TIMS technique on garnet amphibolites with relict eclogitic garnet from the Argentera Massif and eclogites from the Belledonne and Aiguilles Rouges Massifs. They obtained mainly discordant data, whose upper and lower intercepts are of difficult interpretation. In most samples, no age constraints on the HP metamorphism were obtained, but for the Argentera Massif a lower intercept of 424±4 Ma from an amphibolite was proposed as the age of HP metamorphism. Notably, a second mafic rock from the same area returned an upper intercept at ~350 Ma with a meaningless lower intercept. More recently, Carboniferous ages at ~340 Ma were obtained for zircon rims in the Frisson Pl-rich HP granulite and sector zoned zircons of the Pl-poor HP granulite $(340.7 \pm 4.2 \text{ and } 336.3 \pm 4.1 \text{ Ma}, \text{ respectively: Rubatto } et$ al., 2010). Several lines of evidence constrain the formation of the Carboniferous zircon rims to before mylonitization stage C and, possibly, during stage A: 1) the

Explorer HREE depletion in the zircon rims is in line with formation, before or during zircon crystallization, of metamorphic garnet that sequestrated HREE from the reactive bulk rock (Rubatto, 2002); 2) the zircon rims lack a sig-

The Virtual

nificant negative Eu anomaly, which is also absent in the other peak metamorphic minerals such as omphacite, garnet and plagioclase; 3) Ti-in-zircon thermometry indicates temperatures of at least 700-770 °C, which are within that reported for the HP peak of stage A, but generally higher than those of the first retrogression of stage B; 4) the Carboniferous zircon domains show evidence of intense deformation, likely related to the mylonitic stage.





P-T path of the GSV Terrane and metamorphic stages A to D of the Lago Frisson HP granulites, from Ferrando et al. (2008). Phase relations for Al2SiO5 are after Holdaway and Mukhopadhyay (1993) and the wet granite solidus is after Aranovich and Newton (1996).

Metavolcanic rocks

The presence of dacitic to rhyodacitic metavolcanic rocks at Cima Ghiliè-Testa della Rovina and in Val Meris (Fig. 3) is reported by Romain (1982, p.30), Ghiglione (1990), Bierbrauer (1995), Colombo (1996) and Rubatto et al. (2001). The Cima Ghiliè metadacite is undeformed, preserves a porphyritic structure and contains granulitefacies xenoliths. This rock type shows progressive deformation to foliated metadacite (Fig. 7), and transitional contacts to the surrounding migmatites. The undeformed metadacite preserves phenocrysts of quartz, plagioclase, K-feldspar, biotite and altered cordierite (Colombo et al., 1993). The microcrystalline matrix consists of quartz, feldspars and aggregates of biotite + quartz that probably derived from former xenocrysts of granulite-facies orthopyroxene. The metadacite contains pink, euhedral zircons with composite cores, with variable zoning, and oscillatory-zoned overgrowths. The cores yielded concordant U-Pb ages at 560 Ma, 600 Ma and 700 Ma, respectively. The SHRIMP analyses of the oscillatoryzoned overgrowths yielded a mean ${}^{206}Pb/{}^{238}U$ age of 443 ± 3 Ma (Rubatto *et al.*, 2001).

Figure 7. GSV Terrane: metadacite



Slightly foliated metadacite with xenoliths of metamorphic rocks including Grt-Opx granulite. GSV Terrane. Cima Ghilié, upper Gesso della Valletta valley.

The rocks described in the SE GSV Terrane as "pseudoporphyric gneisses" by Malaroda (1991; 1999), though more deformed and recrystallized, appear to be similar to the Ghiliè metadacite both in structure and composition, and are indicated with the same symbol in Fig. 3. Part of the "leptynites", reported especially in the NW of the GSV Terrane in the map of Malaroda et al. (1970), are possibly also derived from recrystallized acid metavolcanics.

Migmatitic paragneisses and migmatitic granitic aneiss

A striking feature of the GSV Terrane is the abundance and variety of migmatites (Malaroda, 1968),



formed from granitic ("anatexites" *Auct.*), pelitic (Fig. 8a) and mafic protoliths. The bulk composition of the "anatexites" is homogeneously granitic (Faure-Muret, 1955; Malaroda & Schiavinato, 1957a; De Pol, 1966; Blasi & Schiavinato 1968). These rocks generally lack structural relicts of the igneous protolith. A notable exception are the "augen anatexite" and "augen embrechite" (Fig. 8b; Malaroda *et al.*, 1970), which are likely

derived from the Late Ordovician porphyritic granitoids that have been extensively documented in the pre-Variscan basement of the External Crystalline Massifs (e.g. Aiguilles Rouges: Bussy & von Raumer, 1993).

Figure 8. GSV Terrane: Migmatites I





(a) Thinly banded paragneiss showing a discordant leucosome with concordant branches developed along an incipient dextral shear zone. GSV Terrane, Laghi della Sella, upper Valle della Meris. (b) Migmatite with K-feldspar porphyroclasts derived from a granitoid of probable Ordovician age. GSV Terrane. Ridge between Valrossa and Valmiana, Valle di Valasco, upper Gesso della Valletta valley.

Figure 9. GSV Terrane: Migmatites II





(a) Thinly banded "anatexite" with both concordant and discordant granitoid leucosomes. GSV Terrane. Path from Rifugio Genova to Colle Fenestrelle, upper Valle della Rovina. (b) "Anatexite" showing cm-thick veins of pinitized cordierite ± quartz corresponding to the latest stages of the anatectic melt crystallisation. GSV Terrane. Upper Valle della Meris. http://virtualexplorer.com.au/

Figure 10. GSV Terrane: Bousset-Valmasque Complex

The Virtual

Explorer



(a) "Granite de la Valmasque". "Agmatite" with angular melanocratic fragments ("enclaves") of hornblendite in a leucogranite matrix. The white patches in the hornblendite are plagioclase-diopside aggregates. GSV Terrane. Upper Valmasque, Pas de la Fous. This outcrop is the same as that described by Faure-Muret (1955, p.100 and Fig.1, Plate XI). (b) Ultramafic fragments are enclosed in an anatectic matrix of granitoid composition. The ultramafite-matrix contact is marked by a continuous rim of radially arranged amphibole, whereas the larger ones still have a serpentinized core. The smaller fragments are completely replaced by amphibole. GSV Terrane. Path between Rifugio Genova and Colle Fenestrelle,upper Valle della Rovina. (c) A larger ultramafic block in a granitoid matrix with sub-idioblastic plagioclase crystals ("perlgneiss"). Path between Rifugio Genova and Colle Fenestrelle, upper Valle della Rovina.

Although a discussion on the protoliths and genesis of the GSV migmatites is outside the scope of this review, we note that leucosomes in the migmatitic paragneiss and migmatitic granitic gneiss have mineral assemblages compatible with the main amphibolite-facies metamorphic stage recorded in the whole GSV Terrane. Specifically, the mineral assemblage observed in metapelites of the upper Val Gesso (Compagnoni et al., 1974; Bierbra-1997)—i.e. quartz-plagioclase uer, 1995; Prever, (An₂₀₋₃₀)-biotite-fibrolitic sillimanite-cordierite with scarce K-feldspar and muscovite—and the relict kyanite and garnet rarely included in plagioclase suggest P-Tconditions of the upper amphibolite-facies (650-700°C; 0.4–0.6 GPa) and decompression from higher P, respectively. In the migmatitic granitic gneiss, more rarely in migmatitic paragneisses, the end products of partial melting are pockets and larger bodies of a fine-grained muscovite-biotite granite ["granito aplitico microgranulare di anatessi" (fine grained aplitic anatectic granite) of Malaroda et al. (1970)] and quartzo-feldspathic pods and dykes that commonly contain large crystals of pinitized cordierite (Fig. 9).

The LP metamorphism and partial melting in the GSV Terrane are not directly dated, but a Late- to Mid-Carboniferous age for migmatization ($\leq 323 \pm 12$ Ma) has been proposed on the basis of a zircon lower intercept age obtained from the Meris eclogite (Rubatto *et al.*, 2001).

The Bousset-Valmasque Complex

The Bousset-Valmasque Complex was named "Entracque-Tenda Granite" by Roccati (1925) and "Granite à enclaves de la Valmasque" by Faure-Muret (1955). It consists of amphibole migmatite and amphibole-bearing anatectic granite that contains enclaves of amphibolite and ultramafics of angular or rounded shape, with size from a few cm to a few meters (Fig. 10). The Complex



comprises two E-W directed subvertical bands separated by the Alpine Fremamorta-Colle del Sabbione shear zone (Fig. 3). The N band extends to a maximum thickness of ~ 5 km between Lago Brocan and the Bousset valley, whereas the S band extends to a maximum thickness of 1 km in the Gelas-Clapier ridge between the Gesso della Barra valley and the upper Valmasque. As noted by Roccati (1925) and Faure-Muret (1955), a noteworthy feature of the Bousset-Valmasque Complex is the great variety of mafic-ultramafic enclaves (including rare eclogite and metagabbro) (Fig. 10a), some of which were studied by Faure-Muret (1955) and Boucarut (1969). The igneous appearance and the lack of deformation of the Valmasque granite suggest either a peak anatectic origin linked to the Late Carboniferous metamorphic event or a post-metamorphic emplacement. However, no geochronological or modern petrological data exist to constrain the age and igneous/metamorphic history of the Bousset-Valmasque Complex.

The Virtual

Explorer

Carboniferous-Permian magmatic rocks

Late Variscan plutonic bodies are widespread in the External Crystalline Massifs of the Alps. The age of emplacement, Mg/(Fe + Mg) ratio, and mafic mineral content allow to recognise two plutonic suites: an earlier, Visean (\sim 330–340 Ma) and Mg-rich suite, and a later suite, mainly Stephanian (\sim 295–305 Ma) and richer in Fe (Debon & Lemmet, 1999).

In the Argentera Massif, rocks belonging to the Visean plutonic suite occur as small bodies and dykes of metamonzonite in the central part of the GSV Terrane, e.g. on the N slope of the Cima dei Gélas and in the upper Valmasque (Figs. 3 and 11). Such rocks were first reported by Roccati (1925) as "pyroxene-bearing porphyritic gneiss" and were later mapped as "biotite-amphibole embrechites" by Malaroda et al. (1970). One of the largest metamonzonite outcrops is exposed near the Muraion glacier, N of Cima dei Gelas. Here the plutonic rock, though showing evidence of incipient migmatization, well preserves the original igneous fabric and mineralogy. The magmatic assemblage consists of cm-sized euhedral crystals of K-feldspar and minor plagioclase in a medium-grained mineral aggregate, which contains relicts of magmatic clinopyroxene that are partly replaced by amphibole and biotite. U-Pb SHRIMP analyses of zircons from this rock yielded a concordant age of 332 ± 3 Ma (Rubatto et al., 2001).

The Stephanian magmatic suite is represented in the GSV Terrane by the Central Granite, a large pluton straddling the boundary between Italy and France in the upper Gesso and Vésubie valleys, and by a minor leucogranite body in the upper Valle Stura (Brondi, 1958; Malaroda et al., 1970, and references therein). The Central Granite is a shallow pluton cutting across the regional metamorphic foliation of country rocks. It was emplaced through magmatic stoping at shallow crustal levels after the Variscan amphibolite-facies metamorphism and anatexis (Boucarut, 1967). Contact metamorphism of the surrounding rocks produced both andalusite and sillimanite (Compagnoni et al., 1974). The Valle Stura leucogranite, on the other hand, is parallel to the regional foliation trend of country rocks, though probably discordant at the outcrop scale.

Figure 11. GSV Terrane: metamonzonite



Deformed metamonzonite where are evident porphyroclasts of the relict igneous K-feldspar and an anatectic dykelet of granitoid composition (centre left), which is cutting across the pre-migmatitic metamorphic foliation . GSV Terrane. Upper Valmasque, just below the Pas de la Fous.

Compositionally, the Central Granite is a biotite-muscovite monzogranite in which four structural and compositional varieties have been distinguished and mapped (Malaroda & Schiavinato, 1957b, Malaroda *et al.*, 1970): 1) medium-grained granite, 2) granite with large crystals of K-feldspar, 3) fine-grained aplitic granite, and 4) finegrained porphyritic granite. Both variety 1 and 3 are locally garnet-bearing. Rounded enclaves of a quartz microdiorite ("enclaves microgrenues plagioclasiques" of Boucarut, 1969) with acicular apatite crystals are common in the central part of the pluton between Lac Nègre and Colle di Fremamorta.

The emplacement age of the Central Granite is constrained by Rb-Sr ages on magmatic muscovites (Ferrara & Malaroda, 1969) at 292 \pm 10 Ma (av. of 4 analyses), whereas a cooling age at ca 296-299 Ma was proposed on the basis of ⁴⁰Ar/³⁹Ar single grains muscovite analyses from the granite (Corsini *et al.*, 2004). The emplacement age of the Valle Stura leucogranite is only constrained by a single Rb/Sr muscovite age from a dyke from Bagni di Vinadio at 299 \pm 10 Ma (Ferrara & Malaroda, 1969).

A probably younger suite of intrusive rocks is represented by quartz-porphyry dykes with large crystals of Kfeldspar exposed on the W ridge of Cima del Dragonet, E of Terme di Valdieri (Franchi 1894; Malaroda *et al.*, 1970). Early Permian-Late Carboniferous basalt-andesite pyroclastics and lava flows are reported from the SE of the GSV Terrane (Romain, 1978), in particular from the Vallon de Figuière Unit (Malaroda ,1999). The widespread swarms of basalt-andesite rocks ["porfiriti anfibolico-plagioclasiche, talora quarzifere" (locally quartzbearing plagioclase-amphibole porphyrites) of Malaroda *et al.* (1970)] that cut across the Central Granite and its country rocks are possibly related to this young volcanism.

Carboniferous sedimentary sequences

Late Carboniferous sediments (conglomerates, sandstones, fossiliferous black schists) are reported from synforms in the GSV gneisses of the upper Vésubie valley (Fig. 3) by Faure-Muret (1955, and references therein) and Haudour *et al.* (1958). Plant fossils indicate a Westphalian D-Stephanian A age (Faure-Muret, 1955; Haudour *et al.*, 1958). Deposition of these sediments marks the final exhumation of the GSV Terrane after the Variscan orogeny.

Unfossiliferous metasediments of presumed Late Carboniferous age (mollieresite Auct.) are described in the area between the Mollières valley and Saint Martin Vésubie (Faure-Muret, 1955; Malaroda *et al.*, 1970). The mollieresite is lithologically similar to the fossiliferous Late Carboniferous—as it consists of conglomerates, sandstones and graphitic schists—but is affected by low-grade metamorphism (Bortolami *et al.*, 1974).

Ferriere-Mollières shear zone

The Ferriere–Mollières shear zone represents the main shear zone cutting the Paleozoic basement in the western portion of the Argentera Massif. It strikes NW–SE and extends from Ferriere (Valle Stura) to the northwest, to Mollières to the southeast. Mylonitic rocks, which are unconformably covered by the Triassic sediments in the Ferriere area, almost continuously crop out along the Ferriere–Mollières shear zone with a thickness ranging from 100 to 1000 m (Fig. 3). These mylonites have been interpreted by Bogdanoff (1986) as a sequence including both metasedimentary rocks ("Micaschistes de La Valetta") and mylonites derived from high-grade metamorphic rocks of both the Tinée and GSV terranes.

In a typical section of the Ferriere-Mollières shear zone between Colle di Stau and Rocco Verde (Valle Stura), mylonites reach a thickness of nearly 1 km (Musumeci & Colombo, 2002). Foliations strike N140-150E and show upright attitude or dip strongly toward northeast or southwest. Mineral lineations are defined by elongated grains of quartz or feldspar, which have a dominant subhorizontal trend, or gently plunge (20°) toward northwest. The mylonites include small bodies of foliated, mica-rich quartzites ["Quarziti del Pebrun" ("Pebrun quartzites") of Malaroda et al., 1970], amphibolites and marbles. In the Rocco Verde section, the Ferriere-Mollières mylonites are interleaved with dm- to m-thick ultramylonitic and phyllonitic layers. In mylonites and phyllonites, foliations dip strongly toward southwest and mineral lineations or slickenside striae are subhorizontal or gently dipping toward northwest. Mylonitic mineral assemblages are indicative of low- to very low-grade metamorphic conditions, whereas the porphyroclasts are derived from high-grade gneiss.

Mylonitic muscovite-bearing leucogranites crop out as subvertical northwest-trending intrusions in the northern portion of the Ferriere–Mollières shear zone. A Rb/Sr muscovite – whole rock age of 327 ± 3 Ma on foliated leucogranite gives a lower limit for the age of mylonitic deformation for which kinematic indicators indicate a dextral sense of shear. This strike-slip tectonics is compatible with the extensional regime that occurred during Carboniferous across western Europe (Musumeci & Colombo, 2002).



Tinée Terrane

The Tinée terrane has been subdivided by Faure-Muret (1955) into three metamorphic formations (Fig. 3): the metasedimentary Varélios-Fougieret, the Anelle-Valabres (see also: Prunac, 1976) and the Rabuons lithological Formations. Part of the Rabuons Formation is exposed on the Italian side of the Argentera Massif in the

Figure 12. Lithologies from Tinée Terrane

upper part of side valleys joining the Stura River from Ferriere to Bagni di Vinadio (Malaroda *et al.*, 1970). Lithologies in this area were grouped into two sequences, the Corborant - and Laroussa Series (Sacchi, 1961; De Pol ,1963), equivalent to the Rabuons and Anelle Formations of the French side, respectively.



(a) Two-micas plagioclase gneiss typical of the Anelle Formation. A biotite-rich foliation with quartz, plagioclase and minor muscovite, sillimanite and garnet wraps around elongated quartz-plagioclase aggregates. Saint Etienne de Tinée, Vallon d'Assuéros, c. 1200 m a.s.l. Faure-Muret collection. Musée d' Histoire Naturelle de Nice, France. Sample 139/46 and Plate II, Fig. 3 of Faure-Muret (1955). (b) Iglière granodioritic to tonalitic gneiss with a biotite foliation wrapping around quartz and feldspar aggregates. Road from Tinée Valley to Roya. Sample 289/45: Faure-Muret collection. Musée d'Histoire Naturelle de Nice, France. (c) Typical Rabuons gneiss, characterized by folded quartz-plagioclase-K-feldspar leucosomes in a quartz-oligoclase-K-feldspar-two micas-sillimanite matrix. Saint Etienne de Tinée, Route de l'Energie, Vallon du Lusernier, 2440 m a.s.l. Faure-Muret collection. Musée d'Histoire Naturelle de Nice, France. Sample 222/45 and Plate IV, Fig. 3 of Faure-Muret (1955). d) Augen gneiss of the Rabuons Formation with a mylonitic foliation. Porphyroclasts of K-feldspar and plagioclase and quartz-feldspar augen enclosed in a closely spaced foliation defined by biotite-rich layers and discontinuous quartz-feldspars layers. Saint Etienne de Tinée, Route de l'Energie, S of Mont Garnet, 2420 m a.s.l. Faure-Muret collection. Naturelle de Nice, France. Sample 248/45 and Plate IV, Fig. 1 of Faure-Muret (1955).

The Anelle-Valabres and Rabuons Formations are less migmatitic than the GSV terrane (Fig. 12). The Rabuons migmatites (Fig. 12c) in particular contain a significantly smaller volume of neosome than the migmatites of the GSV. This neosome forms "augen" or ribbons of K-feldspar and plagioclase or muscovite-bearing granite (Sacchi, 1961; Malaroda, 1968, Tables I, II, III). Both Formations have relicts of more or less retrogressed eclogite (Faure-Muret, 1955; Malaroda *et al.*, 1970). The Anelle-Valabres Formation also includes mafic granulites, coronitic metagabbros (Latouche & Bogdanoff, 1987; Fig. 3) and marbles (Prunac, 1976; Bogdanoff, 1980). In addition to muscovite related to regional metamorphism, the Anelle-Valabres Formation also contains



sillimanite and late kyanite (Fig. 12a). The Rabuons Formation contains muscovite and sillimanite, with scarce kyanite present as relicts in gneisses (Faure-Muret, 1955; Bogdanoff, 1980, 1986) and in quartz veins (Pierrot *et al.*, 1974).

Peculiar amphibole migmatites [amphibole-bearing gneiss and dioritic gneiss ("gneiss anfibolici" and "gneiss dioritici") of Malaroda *et al.*, 1970] that are somewhat similar to the Valmasque Granite, were reported by Franchi (1894, 1933) and Franchi & Stella (1930) in the Ferriere valley, at the NW tip of the Tinée Terrane. Similar migmatites also crop out further to the SE in the upper Bagni di Vinadio valley (Sacchi,1961).

The Tinée terrane is generally lacking large intrusive bodies. Minor intrusions are 1) the Iglière metagranodiorite (Fig. 12b), which was emplaced into the Anelle-Valabres Formation before the Variscan metamorphism and deformations (Bogdanoff & Ploquin 1980), and 2) a small body of aplitic granite and some discordant garnettourmaline-muscovite pegmatite dykes within the Rabuons Formation (Piccoli, 2002).

From his structural work in the upper Tinée valley, Bogdanoff (1980,1986) concluded that the Tinée Terrane records a long deformation history, with complex deformation in Late Proterozoic-Early Paleozoic times (D1). This phase was followed early in the Variscan cycle by HP metamorphism responsible for the formation of HP granulites and eclogites at the expense of basic rocks. A second deformation phase (D2) developed the regional foliation S2. The metamorphic overprint at amphibolitefacies conditions was accompanied by migmatisation. This metamorphism was synchronous with two deformation events at the megascopic scale: nappe stacking and recumbent folding of the Anelle and Rabuons Formations (D3), followed by verticalization of the earlier structures and upright folding at the hectometric scale (D4). A final deformation (D5) consisted of folds of variable style from metric to decametric scale and vertical shear zones, and did not modify significantly the earlier structures.

Time constraints on the evolution of the Tinée terrane are limited to ${}^{40}\text{Ar}/{}^{39}\text{Ar}$ data (Monié & Maluski 1983):

muscovites from the Rabuons Formation yielded and age of 342 ± 7 Ma, the biotites from the Iglière metagranodiorite are as old as ~ 312-316 Ma, and amphiboles and biotites from the Rabuons gneiss were dated at ~ 299 and 283 Ma, respectively.

Mont Blanc - Aiguilles Rouges Massifs

The Mont Blanc, as the Argentera Massif, straddles the boundary between Italy and France in the NW Alps. Elevations in the Mont Blanc Massif are much higher than in the Argentera, culminating at 4810 m in Mont Blanc itself, the highest point in the European continent. This summary is largely based on an extensive compilation of previous works that can be found in von Raumer & Bussy (2004), in Bussy *et al.* (2001), and in the web site at: http://www.unil.ch/webdav/site/igp/shared/stampfli_research/field_trips/field_trip1.pdf

The Mont Blanc Massif (Fig. 13) consists of a complex pile of tectonic units with contrasting peak *P-T* metamorphic conditions. These units are separated by major, steeply dipping NNE-SSW faults and/or mylonitic zones. Most of these tectonic contacts probably formed during the late Variscan strike-slip regime and were reactivated during the Alpine orogeny. One such fault is the *faille de l'Angle* ("Angle fault"; Fig. 13) that separates the Massif in two portions, each one with its sedimentary cover: an internal portion, essentially granitic, to the E, and a more composite external portion to the NW (Epard, 1990).

Unlike the Argentera, the Mont Blanc Massif mainly consists of Variscan granitic rocks, with only minor paragneiss sequences and Ordovician metagranitoids. Its pregranitic history may be better understood in the nearby Aiguilles Rouges Massif, where a polymetamorphic basement is extensively exposed (Fig. 13).

Like the Argentera, the Mont Blanc Massif was only marginally reworked during the Alpine tectonometamorphic cycle, developing low-*T* greenschist-facies mineral assemblages (albite, stilpnomelane, green biotite, epidote, actinolite: von Raumer & Bussy, 2004)



http://virtualexplorer.com.au/





Geotectonic map of the Mont Blanc Massif, from Bussy et al. (2001) and von Raumer & Bussy (2004), modified. Distribution of Carboniferous sedimentary rocks in the Mont Blanc Massif after Gidon (1976).

Polymetamorphic basement and Ordovician metagranitoids

According to the geochronology of its polymetamorphic basement, the Mont Blanc Massif underwent at least three orogenic cycles: Late Precambrian, Ordovician and Variscan (von Raumer *et al.*, 1999).

Magmatic ages of ~ 450 Ma have been obtained from zircons of MORB-like HP mafic rocks (Paquette *et al.*, 1989), and from S-type and I-type calc-alkaline metagranitoids ("Ordovician granitoids": Bussy & von Raumer,

1994). The "Ordovician granitoids" intruded detrital sequences (now paragneisses) of supposedly Late Precambrian to Ordovician age and mainly consisting of sandstones and graywackes, with minor carbonate intercalations and tholeiitic basaltic layers (von Raumer & Bussy, 2004). It has been suggested that flysch-type sediments enriched in Cr and Ni, which are associated to mafic rocks (now eclogites) and ultramafic rocks (Aiguilles Rouges), represent a former accretionary prism (von Raumer, 1998).

The subsequent HP event has not been precisely dated. Although observed in metapelites and paragneisses as relicts of staurolite and kyanite (the latter being sometimes rimmed with reaction coronas of cordierite and spinel), the HP assemblage (1.4 GPa and 700°C) is better preserved in the retrogressed eclogites of Lac Cornu (Aiguilles Rouges, Fig. 13; Liégeois & Duchesne, 1981). An isothermal decompression to ~700 °C and 0.8 GPa and a later amphibolite-facies overprint is preserved in retrogressed eclogites from Val Bérard, Aiguilles Rouges (Fig. 13; Schulz & von Raumer, 1993). Leucocratic rocks associated with some eclogitic boudins have been interpreted as early decompression melting products linked to the exhumation from HP conditions (von Raumer et al., 1996). A preliminary dating of zircons from one of these leucosome yielded an age of ~340 Ma (Bussy & Schaltegger, quoted by Bussy et al., 2001).

The Virtual

Explorer

The paragneisses display a succession of deformation events attributed to the Variscan orogeny, with thrust tectonics (Dobmeier, 1998) and nappe stacking, which led to the development of Barrovian metamorphism (von Raumer et al. 1999). Metapelites of the Aiguilles Rouges experienced a typical Barrovian evolution, recording a clockwise P-T path characterized by sillimanite-bearing assemblages (biotite + plagioclase + quartz + sillimanite \pm garnet \pm cordierite \pm muscovite \pm K-feldspar) and by late stage and alusite found in quartz ± K-feldspar tension gashes (von Raumer, 1984). Metagraywackes and metagranites show evidence of partial melting of variable degree, ascribed to isothermal decompression during exhumation processes. Typical mineral assemblage of these migmatites is plagioclase + quartz + K-feldspar + biotite \pm sillimanite \pm muscovite \pm garnet \pm cordierite. A decompressional event is also documented in polymetamorphic paragneiss collected in the French part of the Mont Blanc Tunnel (Borghi et al., 1987), where an earlier amphibolite-facies mineral assemblage with garnet I, oligoclase, white mica I, is formed before the development of the Variscan regional foliation defined by biotite, white mica II, K-feldspar and garnet II. The thermal peak has been dated at 327 ± 2 Ma by U-Pb analyses on monazites from Lake Emosson micaschists (Fig. 13; Bussy et al., 2000). Monazite from a leucosome vein of a migmatitic graywacke at Lake Emosson yielded a crystallization age of 320 ± 1 Ma, whereas monazite from a migmatitic granite from the Mont Blanc Massif yielded 317 ± 2 Ma (Bussy et al., 2000). These ages are similar to that suggested by Rubatto *et al.* (2001) for migmatization of the Argentera GSV Terrane ($\leq 323 \pm 12$ Ma) on the basis of a zircon lower intercept age obtained for the Meris eclogite.

Greenstone Unit and Visean volcanosedimentary sequence

The SW part of the Aiguilles Rouges Massif (Fig. 13) is characterized by the presence of the so-called "Greenstone Unit" and of low grade metamorphic sediments and volcanics (Dobmeier, 1996; Dobmeier et al., 1999). The Greenstone Unit is composed of greenish, calc-alkaline volcanic rocks of rather low metamorphic grade (von Raumer & Bussy (2004), presumably representing an Early Paleozoic arc environment (Dobmeier et al., 1999), though a Devonian or Silurian age cannot be excluded (von Raumer & Bussy (2004). The low grade metasediments consist of variably deformed pelites, graywackes and sandstones (Bussy et al., 2001), which were metamorphosed at greenschist-facies conditions [chlorite zone, Dobmeier, 1996, 1998)] An Early Carboniferous deposition age was obtained using palynological data (Late Visean acritarchs, Bellière & Streel (1980). Meterthick bands of green metabasalts (Fe-basalts of E-MORB affinity; Dobmeier, 1996) and meta-andesite are found interlayered with the metagraywackes. These rocks possibly record Early Carboniferous transtension linked to the opening of the sedimentary basin and to the 330-340 Ma high-K magmatic pulse.

Carboniferous magmatism

Carboniferous magmatic rocks mainly consist of nonto weakly-metamorphosed intrusions, locally associated to subvolcanic rocks (Fig. 13). Three magmatic pulses are recognized in the Mont Blanc – Aiguilles Rouges area: a) high-K calc-alkaline to shoshonitic, b) peraluminous, and c) metaluminous, ferro-potassic and alkali-calcic (Bussy *et al.*, 2001).

The high-K calc-alkaline to shoshonitic magmatism

The 330-340 Ma high-K calc-alkaline to shoshonitic event that is well known throughout the European Variscan belt is documented by the Pormenaz monzonite (333 \pm 2 Ma; Bussy *et al.*, 2000) and the Montées-Pélissier granite (331 \pm 2 Ma; Bussy *et al.*, 2000), both in the southwestern Aiguilles Rouges.

The Pormenaz monzonite was emplaced at shallow depth, along the transpressive border fault of the Visean

volcano-sedimentary basin. The main facies is porphyritic to equigranular with pink or white cm-sized K-feldspar megacrysts in a dark grey-green amphibole-rich matrix. Euhedral crystals of brown titanite are clearly visible in hand specimen. Metre-sized bodies of durbachite are found as dark-green microgranular magmatic enclaves (von Raumer & Bussy, 2004).

The Virtual

Explorer

Like the Pormenaz monzonite, the Montées-Pélissier granite was emplaced syntectonically along ductile shear zones during the transcurrent stage, and recorded the subsequent vertical movements during cooling (Dobmeier, 1996). It is a foliated fine-grained two-mica monzogranite, which hosts rare biotite-rich restites, mafic microgranular enclaves and biotite-bearing lamprophyres (von Raumer & Bussy, 2004). The lamprophyre-type mafic magmatism is systematically associated to the Mg-K granitoids: the isotopic composition of Sr, Nd and Hf, and the episodic zircon inheritance all point to an essentially high-K lithospheric mantle source, metasomatized during an earlier subduction event, mixed with variable amounts of lower crust material (von Raumer & Bussy, 2004).

The peraluminous magmatism

The second magmatic pulse is represented by peraluminous granites intruded simultaneously into the Aiguilles Rouges and Mont Blanc Massifs at ~ 307 Ma: the Vallorcine and Montenvers granites, and the Fully granodiorite, respectively (Fig. 13).

The Vallorcine granite (Aiguilles-Rouges Massif) is a syntectonic pluton, which intruded at 306.5 ± 1.5 Ma (Bussy et al., 2000) along a steeply dipping, NE-SW trending, dextral strike-slip shear zone. During Late Variscan, the Vallorcine granite was affected along its SE contact by intense low-T ductile shearing subsequently reworked during the Alpine orogeny. The Vallorcine intrusion is a typical S-type granite with Al-rich minerals, such as cordierite, muscovite and andalusite, abundant zircon inheritance and a crustal stable isotope signature (Brändlein et al., 1994). Two facies have been differentiated in the Vallorcine granite (Brändlein et al., 1994): 1) the topographically lower facies is a biotite-rich monzogranite hosting numerous enclaves (up to 30 cm in size), including xenoliths of country gneiss, hornfelses, micaceous restites with sillimanite and hercynite, cordieritebearing leucogranites, and mafic microgranular enclaves; 2) the upper facies is finer grained, with less biotite and almost devoid of enclaves. Aplites and other types of dykes are concentrated within the wall rocks of the upper contact.

The peraluminous Montenvers granite (Mont Blanc Massif) was emplaced syntectonically at 307 ± 3 Ma (Bussy & von Raumer, 1994) as a sheet-like intrusion, similarly to the Vallorcine granite. It is a S-type granite now strongly deformed and often converted to a mylonit-ic leucocratic orthogneiss hosting both microgranular and restitic enclaves.

The Fully granodiorite (Aiguilles-Rouges Massif; Krummenacher, 1959) is a highly heterogeneous, coarsegrained granodiorite of migmatitic appearance (presence of Schlieren and nebulitic structures) and characterized by clots of pinitized cordierite, scattered K-feldspar megacrysts, and numerous small biotite-rich restitic enclaves. Cordierite-bearing leucogranitic dykes and stocks crosscut the main plutonic facies. The Fully granodiorite is a typical peraluminous, anatectic granitoid with a significant restitic component and abundant Al-rich minerals (garnet, cordierite, muscovite, hercynite). It might represent a deeper and less evolved equivalent of the Vallorcine granite. Micaschists, gneisses, marbles and amphibolites are found as dm-long xenoliths. Enclaves of magmatic rocks include microgranular quartz-diorite to granodiorite and angular to rounded pieces of fine- to coarsegrained gabbros up to one meter in size (Bussy et al., 2000). Zircon and monazite, extracted from granodiorite, leucogranite and gabbro, yielded ages of 307 ± 2 Ma for all these rock types (Bussy et al., 2000). Although solid at time of its incorporation into the anatectic mass, the gabbro enclaves are coeval with the acid magmatism, suggesting large-scale dehydration melting of crustal units in close association with mantellic magmas (von Raumer & Bussy, 2004). These field relations have been related to mantellic basic magmatism that was active at the time of crustal anatexis (bimodal magmatism: Bussy et al., 2001).

The metaluminous, ferro-potassic, alkali-calcic magmatism

The last magmatic pulse within the Mont Blanc – Aiguilles Rouges area is represented by the voluminous 303 \pm 2 Ma Mont Blanc granite (Fig. 13), a foliated, porphyritic monzo- to syenogranite with K-feldspar megacrysts and Fe-rich biotite (Marro, 1988; Bussy, 1990). It hosts numerous mafic microgranular enclaves, calc-alkaline



micro-monzodioritic stocks and syn-plutonic dykes of mantellic origin, which record magma mingling processes (Bussy, 1990). The Mont Blanc granite is a metaluminous, ferro-potassic, alkali-calcic intrusion characterized by high K, Y, Zr contents and Fe/Mg ratios, and a low 87 Sr/ 86 Sr initial isotopic ratio of about 0.706 (Bussy *et al.*, 1989). Emplacement of the Mont Blanc granite is the last major magmatic event recorded in the area. However, ash-fall deposits, embedded at different levels of the Salvan Dorénaz basin of the Aiguilles Rouges Massif, have been dated at 295 + 3-4 Ma (Capuzzo & Bussy, 2000). Such ash deposits probably derive from volcanic centers located in the Aar-Gotthard Massifs of Central Alps (Capuzzo & Bussy 2000).

Carboniferous sedimentation

Upper Carboniferous sedimentation is best documented in the Salvan-Dorénaz basin of the Aiguilles Rouges (Fig. 13) where it started at the end of the Westphalian. These coarse-grained clastic sediments formed an alluvial fan system and were deposited on top of the exhumed polymetamorphic basement intruded by Variscan granitoids (Capuzzo, 2000, quoted by Bussy *et al.*, 2001). Subsequent sedimentation of braided, anastomozed and meandering river deposits records a sedimentary evolution in a strike-slip tectonic regime. Volcanic layers within the sediments are dated to the Late Carboniferous (Capuzzo & Bussy, 2000), with an age of 308 ± 3 Ma for basal dacitic flows and of 295 ± 3 Ma for a tuff layer from the upper levels of the basin sequence.

Maures-Tanneron Massifs

Also known as "Provence Crystalline", the Maures (to the SW) and Tanneron (to the NE) Massifs (Fig. 14) were not involved in the Alpine orogeny and preserve a fine example of the South Variscan crust as exposed at the end of the Variscan orogeny. They have been the focus of numerous studies in recent times. Only a brief summary of their geological history is given here to put the geological record preserved in the Argentera and Mont-Blanc-Aiguilles Rouges Massifs in a broader Variscan framework. This summary is based on Toutin-Morin *et al.* (1994) and on the papers by Crévola & Pupin (1994), Elter *et al.* (2004), Bellot (2005), Demoux *et al.* (2008) and Rolland *et al.* (2009), in which the reader will find references to earlier papers.

Maures Massif

The Maures Massif comprises two main tectono-metamorphic blocks, the Western Block and the Eastern Block, which are in tectonic contact along the Grimaud Fault (Fig. 14). The two blocks together display a complete metamorphic zoning, ranging from low greenschistfacies to upper amphibolites-facies conditions and including in the Western Block, from W to E, a chlorite zone, a garnet-chlorite zone, a biotite-staurolite zone, a biotite-kyanite zone, a biotite-muscovite-sillimanite zone, and in the Eastern Block a biotite-sillimanite zone characterized by partial melting (Fig. 14).

The Western Block

The Western Block consists of three metamorphic units: the Western metamorphic Unit, the middle Bormes Unit, and the eastern La Garde-Freinet Unit (Fig. 14).

The Western metamorphic Unit is mainly composed of phyllite and quartzite. Intercalated beds of black schist preserve the well known Llandovery graptolite fauna of Mont Fenouillet, near Hyères (Gueirard *et al.*, 1970), discovered by Schoeller (1938).

The Bormes Unit consists of staurolite-kyanite-garnet two-mica micaschist and two-mica paragneiss, and of the Bormes granitic orthogneiss. Relicts of an aluminous porphyritic granite ("Barral granite"), which was metamorphosed under LP granulite-facies conditions (P =0.67 GPa and $T > 850^{\circ}$ C; Gueirard, 1976), are found in the Bormes orthogneiss. Geochronological data summarized by Bellot (2005) indicate a Precambrian age (~ 550-600 Ma) for the emplacement of the Barral granite. U-Pb dating of monazite from the Bormes orthogneiss yielded an age of 345 ± 3 Ma, interpreted as dating the regional intermediate P metamorphism (Moussavou, 1998). ⁴⁰Ar/³⁹Ar dating of biotite from a sheared orthogneiss yielded ages of 323-328 Ma, that have been interpreted to date the top-to-the NW shear deformation and retrograde low-P metamorphism (Gaubert, 1994, quoted by Bellot, 2005). Metapelites included in the Bormes orthogneiss preserve pre-D1 whiteschist assemblages (1.2-1.6 GPa and 480-550°C) retrogressed to 0.4-0.6 GPa and 600-650°C during shear deformation of the Bormes orthogneiss (Leyreloup et al., 1996, quoted by Bellot, 2005).







Geotectonic map of the Maures and Tanneron Massifs [from Elter et al. (2004) and Rolland et al. (2009), Crévola & Pupin (1994), modified].

The La Garde-Freinet Unit ("Cavalaire Unit" of Bellot, 2005) consists of a wide range of rock types, including acid, mafic, and ultramafic igneous rocks hosted by two mica-garnet-sillimanite micaschist and migmatitic paragneiss. In addition to migmatitic orthogneiss this unit is characterized by an association of amphibolites and "leptynites" (Seyler, 1986) and by the local presence of cordierite in the migmatitic paragneiss. Emplacement of the felsic volcanic protoliths of the "leptynites" is constrained at 548 +15/-7 Ma by U-Pb zircon ages (Innocent *et al.*, 2003). A recent study of the ultramafic rocks (Bellot *et al.*, 2010) shows that they are spinel peridotites and minor garnet–spinel peridotites. In particular, spinel peridotites represent both residual dunites and harzburgites typical of fore-arc upper mantle, and ultramafic cumulates, i.e. dunite adcumulates, harzburgite heteradcumulates and mesocumulates, melagabbro heteradcumulates and amphibole peridotites. The Maures spinel peridotites and melagabbros are interpreted as the lowermost parts of a crustal sequence and minor residual mantle of lithosphere generated in a supra-subduction zone during the Early Paleozoic (Bellot *et al.*, 2010).

The Eastern Block

The Eastern Block ("Cavalières Unit" of Bellot, 2005) is mainly made up of migmatites derived from orthogneiss and biotite-sillimanite paragneiss containing amphibolitized eclogite lenses recording an early HP metamorphism (Fig. 14). Small bodies of biotite- and amphibole-rich orthogneisses are interpreted as primary calc-alkaline granodiorite and diorite, emplaced during the Precambrian (U-Pb zircon ages of 612-630 Ma; Lancelot et al., 1998, quoted by Bellot, 2005). A detailed study of eclogites (Caruba, 1983, Chapt. IX) shows that, even in the best preserved varieties ("amphibolites à grenats auréolés") such as at Cap du Pinet, garnet is the only pristine eclogite phase that survived the subsequent retrogression. A notable exception is the Cavalières eclogite, in which kyanite also survived with garnet, and is surrounded by spectacular corona textures of sapphirine-anorthite symplectite (Bard & Caruba, 1982). The Cavalières kyanite eclogites are inferred to be derived from calc-alkaline gabbro emplaced in a back-arc setting (Buscail et al., 1999 quoted by Bellot, 2005) during Upper Ordovician (zircon U-Pb 452 ± 8 Ma; Lancelot et al., 1998, quoted by Bellot, 2005) and transformed into eclogite during the Lower Silurian (zircon U-Pb 431 ± 4 Ma; Lancelot *et al.*, 1998, quoted by Bellot, 2005).

The Virtual

Explorer

Four generations of synkinematic to postkinematic plutons intruded the central and eastern Maures from 336 to 297 Ma. The oldest intrusions are the Hermitan Granite in Central Maures emplaced at 338 ± 6 Ma and the Reverdit Tonalite in Eastern Maures emplaced at 334 ± 3 Ma (Moussavou, 1998). The largest granite body is represented by the Plan de la Tour Granite of Central Maures, 19 km-long and 5 km-wide (in its northern part), elongated N-S (Fig. 14). The granite is porphyritic-with K-feldspar phenocrysts up to a few cm-long-and hosts dark microgranular "enclaves" or mica-rich inclusions ("enclaves surmicacées") with diffuse rims. Local varieties also contain cordierite crystals a few mm to 1-2 cm of size. The emplacement of the Plan de la Tour Granite is constrained by an U-Pb age of 324 ± 5 Ma (Moussavou, 1998) and by a Rb-Sr age of 313 \pm 10 Ma (Maluski, 1971). The youngest granite intrusion is that of Cap Camarat which was emplaced at c. 300 Ma (Morillon et al., 2000).

The exhumation of the Maures Massif

A detailed ⁴⁰Ar/³⁹Ar geochronological study (22 plateau ages on single grains and bulk samples of amphibole, muscovite and biotite) of metamorphic and magmatic rocks from both sides of the Grimaud Fault (Morillon et al., 2000), shows that migmatites, micaschists, gneisses, pegmatites and granites display muscovite and biotite plateau ages which range from 317.2 ± 1.0 to 322.9 ± 1.7 Ma on the western side of the Grimaud fault, and from 300.2 ± 0.6 to 306.0 ± 2.4 Ma on the eastern side, respectively. In the Western Block, amphiboles yielded plateau ages of 328.1 ± 2.8 and 329.9 ± 2.1 Ma for amphibolites, whereas in the Eastern Block, amphibolite and magmatic bodies contain amphibole that yielded ages ranging from 307.9 ± 1.2 to 317.4 ± 2.4 Ma. These data demonstrate distinct cooling histories in the eastern and western side of the Grimaud Fault. Two distinct periods of fast cooling were recognized, at 320 Ma to the west and at 305-300 Ma to the east of the Grimaud fault, which is identified as a major crustal discontinuity during the late Variscan orogeny.

Post-metamorphic exhumation of the Maures Massif is also constrained by the Late Carboniferous sedimentation in the Plan de la Tour basin. This 16 km-long and 1 km-wide basin is located in the central Maures along the Grimaud fault (Bellot, 2005, and references therein) and is filled with a 400 m thick sequence of conglomerates and arkoses. The base of the sedimentary sequence is dated as Early Stephanian, whereas microgranite dykes emplaced within the basin are of Late Stephanian age (295.4 \pm 2.4 Ma, ⁴⁰Ar/³⁹Ar on biotite: Morillon, 1997, quoted by Bellot, 20 05).

Tanneron Massif

The Tanneron Massif is the 40 x 15 km area of metamorphic rocks and granites sited NE of the Maures Massif, from which it is separated by a belt of Permian sedimentary and volcanic deposits (Fig. 14). The Tanneron Massif is divided by the N-S-trending Joyeuse (to the W) and La Moure (to the E) faults into three domains: the Western, the Central and the Eastern domain.

To the W of the Joyeuse fault (Fig. 14), the Western domain comprises a narrow band of sillimanite-rich migmatitic paragneiss and micaschist forming the basement of the small Late Carboniferous Pennafort basin (G. Crévola in Toutin-Morin *et al.*, 1994; Demoux *et al.*, 2008, and references therein).

To the E of the Joyeuse fault (Fig. 14), the Central domain comprises sillimanite-rich migmatitic gneiss intruded by the granite and tonalite of the Rouet-Prignonet magmatic complex. The Rouet granite, an aluminous porphyritic biotite granite with large crystals of altered cordierite, is comparable to the Plan de la Tour granite of the Maures Massif (Crévola & Pupin, 1994; Onezime et al., 1999). The migmatitic gneisses are sillimanite-rich, occasionally with cordierite, and are intruded by an E-Wtrending tourmaline-bearing leucogranite body (Grime leucogranite) associated with a swarm of leucogranite dykes. More to the E, a thick sequence of orthogneisses and paragneisses, showing increasing partial melting from E to W, and a wide band of blastomylonitic orthogneiss of granodiorite composition (Bois de Bagnols orthogneiss) is found (G. Crévola in Toutin-Morin et al., 1994). The Bois de Bagnols orthogneiss is the basement of the Reyran Late Carboniferous basin, which is filled with 1000 m of sandstone, coal, pelite, and conglomerate dated as Upper Westphalian to Lower Stephanian and hosts intercalations of pyroclastic rhyolite (Basso, 1985, quoted by Bellot, 2005).

The Virtual

Explorer

The Eastern domain, E of the La Moure fault (Fig. 14), is composed of a thick sequence of migmatitic gneiss and of biotite-muscovite-sillimanite-kyanite-garnet micaschist. A migmatitic orthogneiss (Tanneron orthogneiss), intercalated in this series as a N-S-trending bands, locally preserves granitic structure with garnet and cordierite, in spite of widespread transformation into blastomylonitic orthogneiss (Crevola & Pupin, 1994). Amphibolites lenses with eclogite relicts are found in all three domains (Fig. 14). However, even in the best preserved variety ("kelyphitoid eclogite") omphacite is completely replaced by a plagioclase + diopside symplectite (G. Crévola in Toutin-Morin *et al.*, 1994).

Recent isotope dilution U–Pb monazite dating (Demoux *et al.*, 2008) indicates a pre-Variscan history in the Bois de Bagnols orthogneiss of the Central domain, with monazite ages from ~ 440 to 410 Ma. Monazites from a migmatitic paragneiss of the Central domain record a Late Carboniferous HT event at 317 ± 1 Ma (Demoux *et al.*, 2008) In the eastern part, a migmatization event is recorded by monazites from a synkinematic leucogranite body and the Tanneron mylonitic orthogneiss, which yielded ages of 309 ± 5 and 310 ± 2 Ma, respectively (Demoux *et al.*, 2008) Later post-collisional magmatism is recorded by the intrusion of the post-tectonic Rouet granite at 302 ± 4 Ma (Demoux *et al.*, 2008) and of undeformed leucogranite dykes in the Eastern domain, which mark the final stage of the Variscan evolution at 297 ± 5 Ma (Demoux *et al.*, 2008)

Thirty-two ⁴⁰Ar/³⁹Ar plateau ages on single grains of muscovite from metamorphic and magmatic rocks sampled along an east-west transect through the Massif range from 302 ± 2 to 321 ± 2 Ma, and reveal a heterogeneous exhumation that lasted about 20 Ma during the Late Carboniferous (Corsini et al., 2010). In the Eastern Domain, closure of the K-Ar isotopic system is at ~ 311-315 Ma (Corsini et al., 2010). whereas in the Central Domain the K-Ar system closed earlier (~ 317-321 Ma, (Corsini et al., 2010). These cooling paths are interpreted as resulting from differential exhumation processes of distinct crustal blocks controlled by the La Moure fault that separates the two domains (Corsini et al., 2010). In the Western Domain, the ages decrease from ~ 318 to 304 Ma approaching the Rouet granite, which provided the youngest age at 303.6 ± 1.2 Ma (Corsini et al., 2010).

Discussion

The recent geochronological-geochemical studies of Rubatto et al. (2001, 2010) and the petrological study of Ferrando et al. (2008), combined with previous data, well constrain the evolution of the GSV Terrane of the Argentera massif before and during the Variscan orogeny (Fig. 15). On the contrary, the geological history of the Tinée Terrane is less documented, but the presence of eclogites and HP granulites suggests it may have been similar to that of the GSV Terrane. A major difference between the two terranes is that in the Tinée muscovite was stable during the Carboniferous amphibolite-facies event (see below) and the *P*-*T* path ended in the kyanite stability field, not in the cordierite stability field as in the GSV Terrane. Though each massif has its peculiarities and has been studied from somewhat different perspectives, the main geological events recorded in the GSV Terrane of Argentera are also recorded in the Mont Blanc-Aiguilles Rouges and Maures-Tanneron (Table 1).

http://virtualexplorer.com.au/

Figure 15. Argentera Massif: Evolution of the GSV Terrane

The Virtual

Explorer

~ 480 - 460 Ma Intrusion in ortho- and paragneisses of mafic magmas with high degree of crustal contamination

 443 ± 3 Ma Emplacement of dacite sills

340 ± 4 Ma HP granulite-facies metamorphism

> 332 ± 3 Ma Intrusion of monzonite

323 ± 12 Ma Amphibolite-facies metamorphism and anatexis

> 299 - 298 Ma Deposition of Stephanian sediments

~ 295 Ma Intrusion of the Central Granite



The GSV Terrane of the Argentera Massif underwent in the Paleozoic a complex magmatic and metamorphic evolution beginning in the Early Paleozoic (or possibly still in the Late Proterozoic) with the intrusion of large granitoid bodies into a (meta-) sedimentary sequence of graywackes, pelites and rare limestones. In the Maures-Tanneron Massifs, this event is documented by the emplacement of the Barral granite (550-600 Ma) and of the felsic volcanic protoliths of the "leptynites" (~ 548 Ma). However, evidence of this event appears to be lacking in the Mont Blanc-Aiguilles Rouges Massifs (Table 1).

These early granitic intrusions were followed in the Early Ordovician by the emplacement of mafic magmas (now represented by eclogites and HP granulites) in the Argentera (at 480-460 Ma; Fig. 15), in the Maures-Tanneron (at 452 \pm 8 Ma), and in the Mont Blanc-Aiguilles Rouges Massifs (at ~ 450 Ma). In the Argentera Massif the crustal contamination shown by the Laghi del Frisson mafic sequence supports an extensional setting, in agreement with what proposed for other External Crystalline

Massifs (Ménot & Paquette, 1993; Guillot & Ménot, 2009).

A new magmatic cycle, at the Ordovician - Silurian boundary is documented by shallow sheet-like intrusions of dacite with granulitic xenoliths at 443 ± 3 Ma in the GSV Terrane (Fig. 15), by the emplacement of both the S-type and I-type calc-alkaline "Ordovician granitoids" at ~ 440 Ma in the Mont Blanc-Aiguilles Rouges Massifs, and of the granodioritic protolith of the Bois de Bagnols orthogneiss at 440-410 Ma in the Tanneron Massif (Table 1). Granulite xenoliths in the GSV Terrane metadacite are the only preserved evidence of a high-grade metamorphic basement already present in the area during Late Ordovician. Another relic of a LP granulite-facies metamorphism (P = 0.67 GPa and T >850°C;) is reported from the "Barral granite", an aluminous porphyritic metagranite preserved in the "Bormes orthogneiss" (Gueirard, 1976).

These crystalline basements were later subjected to H*P*–H*T* metamorphism with formation, depending from the protolith bulk chemical composition, of either eclogites or H*P*-granulites. *P*-*T*-*t* conditions of this event are well constrained at ~ 1.38 GPa, 735 \pm 15 °C and 340 \pm 4 Ma in the GSV Terrane (Fig. 15). Similar peak metamorphic conditions (1.4 GPa and 700°C) are recorded in the Aiguilles Rouges Massif.

Peak conditions reached by the Laghi del Frisson mafic sequence indicate a geothermal gradient of ~ 16 $^{\circ}C/$ km, compatible with subduction zone environments and, in particular, with the conditions reported by O'Brien (2000) for subduction during continental collision. Relatively high exhumation rates can be inferred for the GSV Terrane from the preservation of peak mineral compositions and the recording of an almost isothermal decompression in the Laghi del Frisson mafic sequence. It is noteworthy that a similar P-T evolution has been reported from the Aiguilles Rouges Massif, where the decompression melting observed around eclogite boudins has been dated at ~ 340 Ma (Table 1). Also the Barrovian amphibolite-facies metamorphism of the Maures "leptynites" at 348 Ma possibly represents decompression following a HP metamorphic event, but this speculation awaits to be confirmed by further studies.



Table 1. Geological evolution of the Argentera, Mont Blanc-Aiguilles Rouges and Maures-Tanneron massifs. References of age data are reported in the text.

| Argentera (GSV Terrane) | Mont Blanc – Aiguilles Rouges | Maures - Tanneron |
|--|---|--|
| Early Paleozoic (possibly Late Pro- terozoic) intrusion of granitoids into a (meta-) sedimentary sequence | | Precambrian (600-550 Ma) emplace- ment of the Barral granite. Emplace- ment at ~ 548 Ma of the felsic volcan- ic protoliths of the Maures "leptyn- ites". |
| Early Ordovician (480-460 Ma) em- placement of the gabbro protoliths of eclogites and HP granulites | Magmatic zircons ages of ~ 450 Ma from eclogitized mafic rocks | Late Ordovician $(452 \pm 8 \text{ Ma})$ emplacement of the calc-alkaline gabbro protolith of the Cavalières kyanite eclogite. |
| Late Ordovician (443 ± 3 Ma) shal- low sheet-like intrusions of dacite with granulite xenoliths | ~ 440 Ma S-type and I-type calc-al- kaline granitoids ("Ordovician gran- itoids") | Emplacement of the granodioritic pro- tolith of the Bois de Bagnols orthog- neiss from Central Tanneron (~ 440-410 Ma). |
| Carboniferous $(340 \pm 4 \text{ Ma})$ meta- morphism at the transition between HP granulite- and eclogite-facies. | Decompression melting, observed around eclogite boudins, at ~ 340 Ma | Barrovian amphibolite-facies meta- morphism of the Maures "leptynites" (~ 348 Ma). |
| Intrusion of dykes and small bodies of monzonite (332 ± 3 Ma). | Mg-K plutonism: Pormenaz monzon- ite (332 ± 2 Ma) and Montées-Péliss- ier granite (331 ± 2 Ma) | Syn-kinematic intrusions of: - Hermi- tan granite (338 ± 6 Ma), Central Maures; - Reverdit tonalite (334 ± 3 Ma), Eastern Maures; - Plan de la Tour granite (324 ± 5 Ma), Central Maures. |
| LP upper amphibolite-facies meta- morphism and anatexis (323 ± 12 Ma) | Amphibolite-facies metamorphism: - 327 ± 2 Ma from Lake Emosson mi- caschists (peak-T age);- 320 ± 1 Ma from a Lake Emosson leucosome (crystallization age);- 317 ± 2 Ma from a Mont Blanc massif migmatitic orthogneiss. | Migmatization in Eastern Maures be- tween 325 ± 8 Ma and 334 ± 3 Ma. Late Carboniferous HT metamor- phism:- 317 ± 1 Ma in migmatitic par- agneiss of the Central Tanneron; - 310 ± 2 Ma in the Tanneron mylonitic or- thogneiss. |
| Emplacement of the Valle Stura leu- cogranite (299 ± 10 Ma) | ~ 307 Ma emplacement of syntecton- ic peraluminous granites (Vallorcine and Montenvers granites), and Fully granodiorite. Calc-alkaline rhyolitic dykes in the Mont Blanc massif | K-Ar cooling ages: - 321-317 Ma, Central Tanneron; - 315-311 Ma, Eastern Tanneron;- 323- 317 Ma, Maures W of the Grimaud fault; - 306-300 E of the Grimaud fault. |
| Emplacement of post-tectonic Cen- tral Granite (299-292 Ma) | Emplacement of post-tectonic metal- uminous, ferro-potassic granitoids (303 ± 2 Ma, Mont Blanc granite). | Post-tectonic intrusion of: - 302 ± 4 Ma Rouet granite, Western Tanneron;- 297 ± 5 Ma leucogranite dykes, East- ern Tanneron; - ~ 300 Ma post-kine- matic Camarat granite, Eastern Maures. |



| Argentera (GSV Terrane) | Mont Blanc – Aiguilles Rouges | Maures - Tanneron |
|---|---|---|
| Deposition on the exhumed migma- tites of Westphalian D-Stephanian A continental sediments. | Deposition of Westphalian D-Stepha- nian A continental sediments in the Salvan-Dorénaz basin, with sub-aer- ial dacitic flows at 308 ± 3 Ma and ash-fall depositsat 295 ± 3 Ma. | Deposition in the Plan de la Tour ba- sin (Maures) of Late Carboniferous sediments cut across by microgranite dykes of Late Stephanian age (295 ± 2 Ma). Deposition in the Reyran basin (Tanneron) of Late Westphalian to Early Stephanian sandstone, coal, pel- ites, and conglomerate. |

Exhumation of the HP rocks was followed—after a few million years—by the emplacement of characteristic Mg-K monzonite and granite. In particular, a limited magmatism of likely extensional nature is represented in the GSV Terrane by the intrusion of small dykes and bodies of K-rich monzonite dated at 332 ± 3 Ma (Fig. 15). A more extensive syn-tectonic magmatism is represented in the Mont Blanc-Aiguilles Rouges Massifs by the Pormenaz monzonite (332 ± 2 Ma) and Montées-Pélissier granite (331 ± 2 Ma), and in the Maures Massif by the the Hermitan granite (338 ± 6 Ma) and Reverdit tonalite (334 ± 3 Ma) (Table 1).

The Virtual

Explorer

Only some 10–20 Ma after the Carboniferous HP metamorphism, an amphibolite-facies metamorphism with extensive development of migmatites occurred not only in the Argentera Massif (323 ± 12 Ma; Fig. 15) but elsewhere in the External Crystalline Massifs (Mont Blanc-Aiguilles Rouges: 327-317 Ma) and Variscan Provence (Maures-Tanneron: 317-310 Ma) (Table 1). The tight succession of HP metamorphism and amphibolite-facies anatexis suggests that, at least in the Argentera Massif, the two stages are part of the same metamorphic cycle, where intense melting occurred upon decompression and advective heat transfer. However, further research is needed to better establish the *P-T* path followed by the GSV and Tinée Terranes during the Carboniferous amphibolite-facies event and its timing.

The Variscan igneous cycle is closed by the emplacement at shallow crustal levels of syntectonic peraluminous granites (Table 1)—the Valle Stura leucogranite (299 \pm 10 Ma) in the GSV Terrane (Fig. 15), the Vallorcine and Montenvers granites (~ 307 Ma) in the Mont Blanc-Aiguilles Rouges Massifs, possibly the Plan de la Tour granite of the Maures massif, which however seems to be older (324 \pm 5 Ma), and of post-tectonic, granites (Table 1), i.e. the Central Granite (cooling ages at 299-292 Ma) in the GSV Terrane (Fig. 15), the Mont Blanc granite $(303 \pm 2 \text{ Ma})$ in the Mont Blanc Massif, the Camarat and Rouet granites (~ 300 Ma) in the Maures-Tanneron.

The final exhumation of the Massifs is marked by the deposition on the crystalline basements of Westphalian D-Stephanian A (306-303 Ma, according to the geological time scale of Gradstein & Ogg, 2004) continental sediments (Fig. 15 and Table 1). Currently available geochronological data seem to suggest that in Argentera the Tinée Terrane may have been exhumed a few My earlier than the GSV Terrane, but this has to be corroborated by further investigations.

Conclusions

Though the main geological events recorded by these Massifs are similar, a few of the events documented in the Mont Blanc-Aiguilles Rouges and Maures-Tanneron have not yet been documented in the Argentera Massif. In particular, the Late Proterozoic emplacement of granitoids in a (meta-) sedimentary sequence documented in the Maures and the Late Ordovician granite plutonism documented in the Mont Blanc Massif, still await to be confirmed by geochronological data. Field observations generally support the occurrence of these magmatic events in the Argentera. On the other hand, the age of the HP metamorphic event documented by eclogites and HP granulites found in all the massifs is well constrained so far only in the GSV Terrane of the Argentera Massif.

A discussion of tectonic models for the Variscan belt of the Western Alps and Provence is outside the scope of this review. However, the recent geochronological data from the Argentera Massif (Rubatto *et al.*, 2010) link together in a single orogenic cycle the HP and amphibolitefacies metamorphisms and suggest that the evolution of the Variscan belt may resemble that of present-day



collisional settings, such as the Himalayan belt. Similarities between the Variscan and the Himalayan orogenies include the conditions of HP granulite-facies metamorphism, and the rapid (within 20 Ma) succession of HP peak metamorphism, fast exhumation and widespread late anatexis. Further petrological, structural and geochronological investigations, however, are needed to explore this hypothesis and properly place the Argentera Massif in the framework of the South Variscan belt.

Acknowledgements

We thank the *Musée d'Histoire Naturelle de Nice* in enabling us to access the Faure-Muret Collection. R.C.

and S.F. gratefully acknowledge the support of the Italian Research Programmes of National Interest (P.R.I.N. Cofin 2004: "Evolution of gondwanian and perigondwanian terranes in the Variscides of Western-Central Alps and Sardinia-Corsica Massif", Scientific Project Coordinator L. Cortesogno).

Journal of the Virtual Explorer, 2010 Volume 36 Paper 3

References

Aranovich, L.Y. & Newton, R.C., 1996. H2O activity in concentrated NaCl solutions at high pressures and temperatures measured by the brucite-periclase equilibrium. Contributions to Mineralogy and Petrology: 125, 200–212.

The Virtual

Explorer

- Bard, J.-P. & Caruba, C. 1982. Texture et minéralogie d'une eclogite à disthène-sapphirine-hypersthène-quartz en inclusion dans les gneiss migmatitiques des Cavalières, massif de Sainte Maxime (Maures orientales, France). Comptes Rendus de l'Académie des Sciences de Paris, 294: 103-106 (in French).
- Bellière, J. & Streel, M., 1980. Roches d'âge viséen supérieur dans le massif des Aiguilles Rouges (Haute Savoie).
 Comptes Rendus de l'Académie des Sciences de Paris, 290: 1341-1343 (in French).
- Bellot, J., 2005. The Palaeozoic evolution of the Maures massif (France) and its potential correlation with other areas of the Variscan belt: a review. In: Carosi, R., Dias, R., Iacopini, D., Rosenbaum G. (eds.): The southern Variscan belt. Journal of the Virtual Explorer, 19, ISSN 1441-8142, 10.3809/jvirtex. 2005.00116
- Bellot, J.-P., Laverne, C. & Bronner, G., 2010. An early Palaeozoic supra-subduction lithosphere in the Variscides: new evidence from the Maures massif. International Journal of Earth Sciences, 99: 473-504. 10.1007/ s00531-009-0416-6
- Bierbrauer, K., 1995. Quantitative Verteilung von Deformation und Strukturen im migmatischen Gesteinen der Mittel/ Unterkruste. 3D- Strukturmodellierung im kristallinen Argentera Massiv in der Externzone der Westalpen. PhD Thesis, University of Würzburg, Germany, 225 pp (in German).
- Bigot-Cormier, F., Sosson, M., Poupeau, G., Stéphan, J.-F. & Labrin, E., 2006. The denudation history of the Argentera Alpine External Crystalline Massif (Western Alps, France– Italy): an overview from the analysis of fission tracks in apatites and zircons. Geodinamica Acta, 19: 455–473.
- Blasi, A. & Schiavinato, G., 1968. Significato petrologico dei noduli a sillimanite e dei noduli a cordierite diffusi nelle anatessiti biotitiche del M. Pelago (Massiccio cristallino dell'Argentera). Bollettino della Società Geologica Italiana, 87: 253-275 (in Italian).
- Bogdanoff, S., 1980. Analyse structurale dans la partie occidentale de l'Argentera-Mercantour (Alpes Maritimes). Thèse, Université de Paris-Sud, Orsay, 316 pp (in French).
- Bogdanoff, S., 1986. Evolution de la partie occidentale du massif cristallin externe de l'Argentera. Place dans l'arc alpin. Géologie de la France, 4: 433-453 (in French).
- Bogdanoff, S. & Ploquin, A., 1980: Les gneiss et migmatites du Massif de l'Argentera (Alpes-Maritimes): apport de deux coupes géochimiques. Bulletin de la Société Géologique de France, ser.7, 22: 353-358 (in French).

- Borghi, A., Gallo, L. M. & Porro, A., 1987. Osservazioni petrografiche nel settore francese del traforo del Monte Bianco. Bollettino del Museo Regionale di Scienze naturali (Torino), 5: 69-96 (in Italian).
- Bortolami, G. & Sacchi, R., 1968. Osservazioni geologicopetrografiche sui medi valloni di S. Anna e Rio Freddo (Massiccio Cristallino dell'Argentera). Memorie della Società Geologica Italiana, 7: 37-64 (in Italian).
- Bortolami, G., Callegari, E. & Gosso, G., 1974. Caratteri metamorfici nella copertura permocarbonifera e nel basamento cristallino dell'Argentera (versante meridionale, St. Martin Vesubie, Francia). Memorie della Società Geologica Italiana, 13: 257-267 (in Italian).
- Boucarut, M., 1967. Structure du granite de l'Argentera et style tectonique de l'ensemble de ce massif (Alpes-Maritimes). Comptes Rendus de l'Académie des Sciences de Paris, 264/D: 1537-1576 (in French).
- Boucarut, M., 1969. Note préliminaire sur les enclaves des massifs granitiques de l'Argentera-Mercantour (A.M.-France) en relation avec les conclusions de J. Didier sur les enclaves des massifs granitiques. Schweizerische mineralogische und petrographische Mitteilungen, 49: 77-96 (in French).
- Brändlein, P., Nollau, G., Sharp, Z. & von Raumer, J. F., 1994. Petrography and geochemistry of the Vallorcine-Granite (Aiguilles Rouges Massiv, Western Alps). Schweizerische mineralogische und petrographische Mitteilungen, 74: 227-243.
- Brondi, A., 1958. Studio geologico-petrografico del massiccio granitico dell'Argentera-Mercantour. Studi e Ricerche della Divisione Geomineraria del C.N.R.N., 1: 187-238 (in Italian).
- Bussy, F., 1990. Pétrogenèse des enclaves micro-grenues associées aux granitoïdes calco-alcalins: exemple des massifs varisque du Mont-Blanc (Alpes occidentales) et miocène du Monte Capanne (Ile d'Elbe, Italie). Mémoires de Géologie (Lausanne), 7: 1-309 (in French).
- Bussy, F. & von Raumer, J. F., 1993. U-Pb dating of Paleozoic event in the Mont-Blanc crystalline massif, Western Alps. Terra Nova, 1: 382-383
- Bussy, F. & von Raumer, J. F., 1994. U-Pb geochronology of Palaeozoic magmatic events in the Mont-Blanc Crystalline Massif, Western Alps. Schweizerische mineralogische und petrographische Mitteilungen, 74: 514-515.
- Bussy, F., Schaltegger, U. & Marro, C., 1989. The age of the Mont-Blanc granite (Western Alps): a heterogeneous isotopic system dated by Rb-Sr whole rock determination on its microgranular enclaves. Schweizerische mineralogische und petrographische Mitteilungen, 69: 3-13.



http://virtualexplorer.com.au/

- Bussy, F., Hernandez, J. & von Raumer J. F., 2000. Bimodal magmatism as a consequence of the post-collisional readjustment of the thickened variscan continental lithosphere (Aiguilles Rouges/Mont-Blanc massifs, western Alps). Transactions of the Royal Society of Edinburgh: Earth Sciences, 91: 221-233.
- Bussy, F, von Raumer J. F. & Capuzzo, N., 2001. Mont Blanc — Aiguilles Rouges Massifs — (External Massifs). In G.M. Stampfli (ed.): Geology of the western Swiss Alps, a guidebook. Field-Trip 1. Mémoires de Géologie (Lausanne), 36: 53-85. Also at: www.unil.ch/webdav/site/igp/shared/ stampfli_research/.../field_trip1.pd
- Calisesi, E. 1971. Studio petrografico delle lenti di marmo nei terreni cristallini dell'Argentera (Alpi Marittime). Unpubl. MSc Thesis, University of Torino (in Italian).
- Capuzzo, N. & Bussy, F., 2000. High-precision dating and origin of synsedimentary volcanism in the Late Carboniferous Salvan-Dorénaz basin (Aiguilles-Rouges massif, Western Alps). Schweizerische mineralogische und petrographische Mitteilungen, 80: 147-167.
- Caruba, C. 1983. Nouvelles données pétrographiques, minéralogiques et géochimiques sur le massif métamorphique hercynien des Maures (Var, France): comparaison avec les segments varisques voisins et essais d'interprétation géotectonique. PhD thesis, University of Nice, France, 359 pp (in French).
- Colombo, F., 1996. Evoluzione tettonico-metamorfica del Complesso Malinvern-Argentera (Massiccio Cristallino dell'Argentera, Alpi Marittime, Italia). PhD Thesis, University of Turin, Italy, 125 pp (in Italian).
- Colombo, F., Compagnoni, R. & Lombardo, B., 1994. Le rocce eclogitiche dei Laghi del Frisson (Argentera sud-orientale, Alpi Marittime). Atti Ticinensi di Scienze della Terra (Pavia), Serie Speciale 1: 75–82 (in Italian).
- Colombo, F., Ghiglione, G. & Compagnoni, R., 1993. Relitti di porfidi granitici a xenoliti granulitici nelle migmatiti dell'Argentera (Alpi Marittime). Plinius, 10: 113-116. (in Italian).
- Compagnoni, R., Lombardo, B. & Prato, R., 1974. Andalousite et sillimanite aux contacts du granite central de l'Argentera (Alpes Maritimes): Rendiconti della Società Italiana di Mineralogia e Petrologia, 30: 31-54 (in French).
- Corsini, M., Ruffet, G. & Caby, R., 2004. Alpine and latehercynian geochronological constraints in the Argentera Massif (Western Alps). Eclogae geologicae Helvetiae, 97: 3– 15.
- Corsini, M., Bosse, V., Féraud, G., Demoux, A. & Crévola, G., 2010. Exhumation processes during post-collisional stage in the Variscan belt revealed by detailed 40Ar/39Ar study (Tanneron Massif, SE France). International Journal of Earth Sciences, 99: 327-341.

- Crévola, G. & Pupin, J. P., 1994. Crystalline Provence: structure and Variscan evolution. In: Keppie, J.D. (ed.): Pre-Mesozoic Geology in France and Related Areas. IGCP Project 233: Terranes in the Circum-Atlantic Paleozoic Orogens. Springer-Verlag, 514 pp.
- Debon, F. & Lemmet, M., 1999: Evolution of Mg/Fe ratios in late Variscan plutonic rocks from the external crystalline massifs of the Alps (France, Italy, Switzerland). Journal of Petrology, 40: 1151–1185.
- Demoux, A., Scharer, U. & Corsini, M., 2008. Variscan evolution of the Tanneron massif, SE France, examined through U–Pb monazite ages. Journal of the Geological Society, 165: 467–478.
- De Pol, C., 1963. Migmatiti e metamorfiti della regione M. Corborant-M. Laroussa (Massiccio dell'Argentera). Rendiconti della Società Mineralogica Italiana, 19: 79-98 (in Italian).
- De Pol, C., 1966. Le anatessiti del M. Ischietto (Massiccio dell'Argentera). Rendiconti dell'Istituto Lombardo di Scienze e Lettere, 100: 77-98 (in Italian).
- Dobmeier, C., 1996. Geodynamische Entwicklung des südwestlichen Aiguilles-Rouges-Massivs (Westalpen, Frankreich). Mémoires de Géologie (Lausanne), 29: 1-198 (in German).
- Dobmeier, C., 1998. Variscan P-T deformation paths from the southwestern Aiguilles Rouges massif (External massif, western Alps) and their implication for its tectonic evolution. Geologische Rundschau, 87: 107-123. 10.1007/ s005310050193
- Dobmeier, C., Pfeifer, H. R. & von Raumer, J. F., 1999. The newly defined "Greenstone Unit" of the Aiguilles-Rouges massif (western Alps); remnant of an early Palaeozoic oceanic island-arc? Schweizerische mineralogische und petrographische Mitteilungen, 79: 263-276.
- Elter, F. M., Corsi, B., Cricca, P. & Muzio, G., 2004. The southwestern Alpine foreland: correlation between two sectors of the Variscan chain belonging to "stable Europe": Sardinia -Corsica and Maures Massif (south-eastern France). Geodinamica Acta, 17: 31-40. 10.3166/ga.17.31-40
- Epard, J. L., 1990. La nappe de Morcles au sud-ouest du Mont-Blanc. Mémoires de Géologie (Lausanne), 8: 165 pp (in French).
- Faure-Muret, A., 1955. Etudes géologiques sur le massif de l'Argentera-Mercantour et ses enveloppes sédimentaires. Mémoires pour servir à l'explication de la Carte géologique détaillée de la France, Paris, Imprimerie Nationale, France. 336 pp (in French), with «Esquisse Géologique du Massif de l'Argentera-Mercantour et de sa Bordure Sédimentaire (Versant français)» at the 1/100.000 scale.

Ferrando, S., Lombardo, B., Compagnoni, R., 2008. Metamorphic history of HP mafic granulites from the Gesso-Stura Terrain (Argentera Massif, Western Alps, Italy). European Journal of Mineralogy, 20: 777–790.

The Virtual

Explorer

- Ferrara, G. & Malaroda, R., 1969. Radiometric age of granitic rocks from the Argentera Massif (Maritime Alps). Bollettino della Società Geologica Italiana, 88: 311–320.
- Franchi, S., 1894. Relazione sui principali risultati del rilevamento geologico nelle Alpi Marittime eseguito nelle campagne 1891-92-93. Bollettino del Regio Comitato Geologico, 25: 231-258 (in Italian).
- Franchi, S., 1933. Carta Geologica d'Italia 1:100.000: Foglio 90 (Demonte). 1a ed., Regio Ufficio Geologico. Stabilimento Tipografico L. Salomone, Roma.
- Franchi, S. & Stella, A., 1930. Carta Geologica d'Italia 1:100.000: Foglio 78-79 (Argentera-Dronero). 1a ed., Regio Ufficio Geologico. Stabilimento Tipografico L. Salomone, Roma.
- Ghiglione, G., 1990. Studio geologico dell'alta Valle della Rovina (Massiccio dell'Argentera). Unpubl. MSc Thesis, University of Turin, Italy, 393 pp (in Italian).
- Gidon, M., 1976. Carte géologique simplifié des Alpes Occidentales du Léman à Digne. Feuille Nord. Editions Didier-Richard and Bureau de Recherches Géologiques et Minières.
- Gradstein, F. M., Ogg, J.G., Smith, A. G. et al., 2004. A Geologic Time Scale 2004. Cambridge University Press. Cambridge, U.K., 610 pp.
- Gueirard, S. 1976. Le granite de Barral, un nouveau témoin des conditions sévères de métamorphisme qui ont affecté la région occidentale du Massif des Maures. Comptes Rendus de l'Académie des Sciences de Paris, 283: 455-457 (in French).
- Gueirard, S., Waterlot, G., Gherzi, A. & Samat, M., 1970. Sur l'âge Llandovérien supérieur à Tarannonien inférieur des schistes à Graptolites du Fenouillet, massif des Maures (Var). Bulletin de la Société géologique de France, serie 7, 12: 195-199 (in French).
- Guillot, S. & Ménot, R. P., 2009. Paleozoic evolution of the External Crystalline Massifs of the Western Alps. Comptes Rendus Geosciences, 341: 253–265.
- Haudour, J., Sarrot-Reynauld, J. & Vernet, J., 1958. Observations nouvelles sur le Houiller et la Mollièrésite dans le massif du Mercantour et comparaison avec le Houiller du dôme de la Mure (Isère). Comptes Rendus de l'Académie des Sciences de Paris, 247/D: 716-719 (in French).
- Holdaway, M.J. & Mukhopadhyay, B., 1993. A re-evaluation of the stability relations of andalusite: thermochemical data and phase diagram for the aluminium silicates. American Mineralogist, 78: 298-315.

- Innocent, C., Michard, A., Guerrot, C. & Hamelin B., 2003. Datation U-Pb sur zircons à 548 Ma de leptynites des Maures centrales. Signification géodynamique des complexes leptyno-amphibolitiques de l'Europe varisque. Bulletin de la Société géologique de France, 174: 585–594 (in French).
- Jung, J. & Roques, M., 1952. Introduction à l'étude zonéographique des formations cristallophylliennes. Bulletin du Service de la Carte Géologique de France, 235:1-62 (in French).
- Krummenacher, D., 1959. Le cristallin de la région de Fully (Valais). Schweizerische mineralogische und petrographische Mitteilungen, 39: 151-266.
- Latouche, L. & Bogdanoff, S., 1987. Evolution précoce du massif de l'Argentera: apport des eclogites et des granulites. Géologie Alpine (Grenoble), 63: 151-164 (in French).
- Liégeois, J. P. & Duchesne, J. C., 1981. The Lac Cornu retrograded eclogites (Aiguilles Rouges Massif, Western Alps, France): evidence of crustal origin and metasomatic alteration. Lithos, 14: 35-48.
- Malaroda, R., 1968. Atlante delle facies petrografiche del Massiccio Cristallino dell'Argentera: Memorie della Società Geologica Italiana, 7: 29-36 (in Italian).
- Malaroda, R., 1991. Gli gneiss-pseudoporfido, particolare tipo litologico dell'Argentera meridionale. Gli affioramenti nel bacino della Beonia (Bieugne) (Alpes-Maritimes, Francia). Atti della Accademia Nazionale dei Lincei (Roma), ser.9a, 1: 129-152 (in Italian).
- Malaroda, R., 1999. L'Argentera meridionale -Memoria illustrativa della "Geological Map of Southern Argentera Massif (Maritime Alps). Memorie di Scienze Geologiche (Padova), 51: 241-331 (in Italian).
- Malaroda, R., Carraro, F., Dal Piaz, G. V., Franceschetti, B., Sturani, C. & Zanella, E., 1970. Carta geologica del Massiccio dell'Argentera alla scala 1:50.000 e note illustrative. Memorie della Società Geologica Italiana, 9: 557–663 (in Italian).
- Malaroda, R. & Schiavinato, G., 1957a. Le anatessiti dell'Argentera. Rendiconti della Società Mineralogica Italiana, 14: 249-274 (in Italian).
- Malaroda, R. & Schiavinato, G., 1957b. Osservazioni preliminari sui fenomeni di anatessi nel settore italiano del massiccio dell'Argentera. Bollettino della Società Geologica Italiana, 76: 323-343 (in Italian).
- Maluski, H., 1971. Etude 87Rb/87Sr des minéraux des gneiss de Bormes (Maures, France). Comptes Rendus de l'Académie des Sciences de Paris, 273: 1470-1473 (in French).



Marro, C., 1988. Organisation géochimique et intrusion du granite du Mont Blanc et de deux leucogranites. Schweizerische mineralogische und petrographische Mitteilungen, 68: 521-529 (in French).

The Virtual

Explorer

- Matte, P., 2001. The Variscan collage and orogeny (480–290 Ma) and the tectonic definition of the Armorica microplate: a review. Terra Nova, 13: 122–128.
- Ménot, R. P. & Paquette, J. L., 1993. Geodynamical significance of basic and bimodal magmatism in the external domains. In: von Raumer, J. F., Neubauer, F. (eds.), Pre-Mesozoic Geology in the Alps. Springer, Berlin, Heidelberg, 241–254.
- Monié, P. & Maluski, H., 1983. Données géochronologiques 39Ar–40Ar sur le socle antépermien du massif de l'Argentera-Mercantour (Alpes-Maritimes, France). Bulletin de la Société géologique de France, 7/25: 247–257 (in French).
- Morillon, A.-C., Féraud, G., Sosson, M., Ruffet, G., Crévola, G. & Lerouge, G., 2000. Diachronous cooling on both sides of a major strike slip fault in the Variscan Maures Massif (south-east France), as deduced from a detailed 40Ar/39Ar study. Tectonophysics, 321: 103-126.
- Moussavou, M. 1998. Contribution à l'histoire thermotectonique varisque du massif des Maures par la typologie du zircon et la géochronologie U/Pb sur minéraux accessoires. Ph.D. thesis, Univ. Montpellier 2, France, 187 pp (in French).
- Musumeci, G. & Colombo, F., 2002. Late Visean mylonitic granitoids in the Argentera Massif (Western Alps, Italy): age and kinematic constraints on the Ferriere-Mollières shear zone. Comptes Rendus de l'Académie des Sciences de Paris, 334: 213-220.
- O'Brien, P.J., 2000. The fundamental Variscan problem: hightemperature metamorphism at different depths and highpressure metamorphism at different temperatures. In: Franke,W., Haak, V., Oncken, O., Tanner, D. (eds.), Orogenic Processes: Quantification and Modelling in the Variscan Belt. The Geological Society, London, 369–386.
- Onezime, J., Faure, M. & Crévola, G., 1999. Petro-structural analysis of the Rouet-Plan-de-la-Tour granitic complex (Maures and Tanneron massifs, Var, France). Comptes Rendus de l'Academie des Sciences, Series IIA, Earth and Planetary Science, 328: 773-779.
- Paquette, J.-L., Ménot, R.-P. & Peucat, J.-J., 1989. REE, Sm-Nd and U–Pb zircon study of eclogites from the Alpine External Massifs (Western Alps): evidence for crustal contamination. Earth and Planetary Science Letters, 96: 181–198.
- Piccoli, G.C., 2002. Minerali delle Alpi Marittime e Cozie meridionali,. Provincia di Cuneo. Associazione Amici del Museo Federico Eusebio (Alba), Savigliano, L'Artistica, 366 pp (in Italian).

- Pierrot R., Picot P., Féraud J. & Vernet J., 1974. Inventaire minéralogique de la France n°4. Alpes Maritimes. Editions du B.R.G.M., Paris, 167 pp.
- Prever, C., 1997. Studio geologico-petrografico dei Valloni d'Orgials, Rio Freddo e dell'alta Valle di Valasco (Massiccio Cristallino dell'Argentera). Unpubl. MSc Thesis, University of Torino, 210 pp (in Italian)
- Prunac M., 1976. Analyse structurale dans les socle de l'Argentera – Mercantour (Région d'Isola: alpes Maritimes). Thèse 3ème cycle, University of Paris Sud, Centre d'Orsay, 123pp.
- Roccati, A., 1925. Il Massiccio Cristallino delle Alpi Marittime. Studio litologico-mineralogico. Bollettino del Club Alpino Italiano., 42/75: 189-241 (in Italian).
- Rolland, Y., Corsini, M. & Demoux A., 2009. Metamorphic and structural evolution of the Maures-Tanneron massif (SE Variscan chain): evidence of doming along a transpressional margin. Bulletin de la Société géologique de France, 180: 217-230.
- Romain, J., 1978. Etude pétrographique et structurale de la bordure sud-occidentale du massif de l'Argentera, de St-Martin-Vésubie à la Cime du Diable. Thèse 3ème cycle, Université de Nice, 300 pp. (in French).
- Romain, J., 1982. Le Massif du Mercantour. Guide géologique. Nice, Ed. Serre, 106 pp (in French).
- Rubatto, D., 2002. Zircon trace element geochemistry: distribution coefficients and the link between U–Pb ages and metamorphism. Chemical Geology, 184: 123–138.
- Rubatto, D., Ferrando, S., Compagnoni, R. & Lombardo, B., 2010. Carboniferous high pressure metamorphism of Ordovician protoliths in the Argentera Massif (Italy), Southern European Variscan belt. Lithos, 116: 65–76.
- Rubatto, D., Schaltegger, U., Lombardo, B., Colombo, F. & Compagnoni, R., 2001. Complex Paleozoic magmatic and metamorphic evolution in the Argentera Massif (Western Alps) resolved with U–Pb dating. Schweizerische mineralogische und petrographische Mitteilungen, 81: 213– 228.
- Sacchi, R., 1961. Notizie geologiche sulla dorsale M. Sejta-M. Corborant (Massiccio dell'Argentera): Atti della Società Italiana di Scienze Naturali, 100: 397-406 (in Italian).
- Schoeller, H., 1938. Sur la présence de Graptolites dans les schistes métamorphiques du massif des Maures. Comptes Rendus Sommaires de la Société Géologique de France, 8: 147-148 (in French).
- Schulz, B. & von Raumer, J. F., 1993. Syndeformational uplift of Variscan high-pressure rocks (Col de Bérard, Aiguilles Rouges Massif, Western Alps). Zeitschrift der deutschen geologischen Gesellschaft, 144: 104-120.



- Stampfli, G.M., von Raumer, J.F. & Borel, G.D., 2002. Paleozoic evolution of pre-Variscan terranes: from Gondwana to the Variscan collision. Geological Society of America Special Paper, 364: 263–280.
- Toutin-Morin, N., Bonijoly, D., Brocard, C., Broutin, J., Crévola, G., Dardeau, G., Dubar, M., Féraud, J., Giraud, J.D., Godefroy, P., Laville, P. & Meinesz, A., 1994. Notice explicative, Carte géologique de France (1/50000), feuille Fréjus-Cannes (1024). Orléans: BRGM, 187 pp.
- von Raumer, J. F., 1984. The external massifs, relics of variscan basement in the Alps. Geologische Rundschau, 73: 1-31. 10.1007/BF01820358
- von Raumer, J. F., 1998. The Palaeozoic evolution in the Alps: from Gondwana to Pangea. Geologische Rundschau, 87: 407-435. 10.1007/s005310050219
- von Raumer, J. F., Abrecht, J., Bussy, F., Lombardo, B., Ménot, R. P. & Schaltegger, U., 1999. The Palaeozoic metamorphic evolution of the Alpine External Massifs. Schweizerische mineralogische und petrographische Mitteilungen, 79: 5-22.

- von Raumer, J. F. & Bussy, F., 2004. Mont Blanc and Aiguilles Rouges: geology of their polymetamorphic basement (External Massifs, Western Alps, France–Switzerland). Mémoires de Géologie (Lausanne), 42: 1–204.
- von Raumer, J. F., Bussy, F. & Stampfli, G. M., 2009. The Variscan evolution in the External Massifs of the Alps and place in their Variscan framework. Comptes Rendus Geosciences, 341: 239–252.
- von Raumer, J. F., Bussy F.A. & Sharp, Z. D., 1996. Lac Cornu revisited: the evolution from lower to upper crust (Aiguilles Rouges Massif, Western Alps). Schweizerische mineralogische und petrographische Mitteilungen, 76: 120-121.